

David Pond

WATER QUALITY REPORT

2021



30 Mile River Watershed Association
P.O. Box 132
Mount Vernon, ME
www.30mileriver.org

2021 David Pond Water Quality Report

TABLE OF CONTENTS

2021 Water Quality Summary	1
Overview	2
Water Quality Monitoring in 2021	2
Secchi Disk Transparency (Water Clarity)	3
Dissolved Oxygen and Temperature	4
Total Phosphorus (TP)	6
Chlorophyll (Chl-a)	7
Discussion	8
Next Steps.....	8

TABLE OF FIGURES


Figure 1. David Pond Monitoring Stations, Maine DEP.....	2
Figure 2. 2021 Secchi Disk Transparency, Station 1	3
Figure 3. Historical Secchi Disk Transparency, Station 1, 1982-2021.....	4
Figure 4. 2021 Dissolved Oxygen and Temperature Profiles, Station 1	5
Figure 5. Annual Average Total Phosphorus data collected 1982-2021, Station 1.	7
Figure 6. Annual average and peak chlorophyll-a, station 1, 1982-2021	7

2021 Water Quality Summary¹

Monitoring on David Pond occurred on 10 dates between May and September 2021 by Whitney Baker of 30 Mile River Watershed Association (30 Mile) and local volunteers.


Water clarity readings in 2021 ranged from 5.66 meters (June 30th) to 6.77 meters (September 22nd) with an annual average of 6.33 meters. 12 readings were collected in 2021 in total.

Water Clarity (m)	
2021 Water Clarity Average	6.3
Historical SDT Average	5.7
Maine Lakes SDT Average	4.8




Five (5) samples were collected and analyzed for **Total Phosphorus**. Laboratory results ranged from 5 ppb (parts per billion) to 8 ppb with an average of 7 ppb.

Total Phosphorus (ppb)	
2021 TP Average	7
Historical TP Average	7
Maine Lakes TP Average	12



Chlorophyll was measured five (5) times in 2021. Results ranged from 2 ppb to 4 ppb with an annual average of 3 ppb.

Chlorophyll-a (ppb)	
2021 Chl-a Average	2.7
2021 Peak Chl-a	4.0
Historical Chl-a Average	3.5
Maine Lakes Chl-a Average	5.4



Nine (9) **Dissolved Oxygen (DO)** profiles were collected in 2021. Anoxia (DO <2 ppm) was first encountered in deep waters at a depth of 11 meters in early June, but this zone of anoxia grew to include all waters 7 meters and deeper by mid-July and persisted through the month of September.

¹ Scale bars illustrate the range of data collected for each parameter over the historical monitoring record for general comparison with the 2021 monitoring results. The blue end represents the historical minimum (best), and the red end represents the historical maximum (worst) of all monitoring data collected.

Overview

David Pond is located in the town of Fayette in Kennebec County, Maine and has a direct watershed area of just over 2 square miles, and an indirect upstream watershed of approximately 2 square miles, which includes the drainages of Tilton and Basin Ponds. David Pond has a single outlet located at the north end of the northern basin that crosses Sandy River Road in Chesterville and continues into Parker Pond.

David Pond is a relatively shallow lake with a maximum depth of 11 m (37 ft) and an average depth of just 3m (10 ft). The Pond has a surface area covering approximately 300 acres. David Pond does not have a public boat launch.




Figure 1. David Pond Monitoring Stations, Maine DEP.

Water Quality Monitoring in 2021

Water quality monitoring on David Pond takes place at the deepest spot in the lake (Maine DEP Station 2), also known as the “deep spot”, located in the south basin. Station 2 is just over 11 meters (37 ft) deep (Figure 1). Monitoring in 2021 was completed by Whitney Baker of 30 Mile River Watershed Association (30 Mile) and local volunteers. A special thanks to our 2021 water quality volunteers on David Pond:



2021 water quality volunteer, Greg Cauldwell.

Greg Cauldwell
Pat & Linda Stanton

Water quality data was collected on 10 dates between May and September. Parameters include Secchi disk transparency, dissolved oxygen and temperature, phosphorus, chlorophyll, and advanced chemistry parameters (pH, Alkalinity, Color, and Conductivity).

Secchi Disk Transparency (Water Clarity)

Secchi disk transparency (SDT) is an indicator of water clarity. To measure water clarity, a black and white disk is lowered into the water and the reading is taken at the depth at which it is no longer visible. Factors that affect water clarity include algal growth, zooplankton densities, natural water color, and suspended silt or sediment particles.

Water Clarity (m)	
2021 Water Clarity Average	6.3
Historical SDT Average	5.7
Maine Lakes SDT Average	4.8

Water clarity readings in 2021 ranged from 5.66 meters (June 30th) to 6.77 meters (September 22nd) with an annual average of 6.33 meters. 12 total readings were collected over 10 monitoring days in 2021 (Figure 2).




Figure 2. 2021 Secchi Disk Transparency, Station 1

SDT data has been collected during 17 years throughout the historical monitoring period spanning the past 39 years. SDT readings in David Pond have ranged from 4.3 m (2009 and 2012) to 8.0 m (2020) with a historical average of 5.7 m (Figure 3).




Figure 3. Historical Secchi Disk Transparency, Station 1, 1982-2021

Dissolved Oxygen and Temperature

Dissolved oxygen (DO) is a critical indicator of the health of the lake system. DO is produced through photosynthesis, consumed during respiration and decomposition, and is influenced by wind, wave action, weather events, and lake productivity. A good supply of oxygen is essential for fish and other aquatic species, with most fish species requiring a DO concentration of 5 ppm (parts per million) or more. As lakes become more biologically productive in the summer, oxygen can decline as decomposition occurs in deep areas of the lake. Loss of oxygen may indicate a stressed and changing ecosystem. Understanding the pattern and extent of oxygen loss in deep waters is important to understanding changes between the years and throughout a single season, and is particular concerning for David Pond because it may be more vulnerable to internal phosphorus loading due to its unique sediment chemistry.²

As lake water is warmed during the summer, lakes will form three distinct temperature layers. There is a warm layer at the surface (epilimnion), a thin transitional layer (metalimnion), and a deep cold layer (hypolimnion) that becomes isolated from the surface and oxygen resupply. In David Pond, severe oxygen loss in the hypolimnion typically occurs in July through September each year.

²David Pond appears on Maine DEP's list of "Threatened Lakes" on the NPS Priority Watersheds List (https://www.maine.gov/dep/land/watershed/nps_priority_list/NPS%20Priority%20List%20-%20Lakes20.pdf) due to its sediment chemistry. Sediment results from samples collected from the bottom of David Pond suggest that the lake is more vulnerable to internal phosphorus loading, a phenomenon that can occur when deep waters become anoxic (DO loss <2 ppm) resulting in phosphorus release from the bottom sediments exposed to anoxic waters.

DO <5 ppm was documented in every profile collected in 2021 and was first encountered in deep waters only (10 meters and deeper) in May, but grew to include all waters 6 meters and deeper by early-August, and persisted through the end of September at this depth. Severe oxygen loss (DO <2 ppm, a.k.a. “anoxia”) was first documented at a depth of 11 meters in early June, but this zone of anoxic water grew to include waters 7 meters and deeper in mid-July, and persisted at this depth through the end of September (Figure 4).

Though oxygen loss in the deep areas of David Pond is not uncommon in recent years, historical data is limited with DO data collection starting somewhat recently in 2013. David Pond supports a warm water fishery (large and smallmouth bass, and chain pickerel) due to the oxygen deficiency in the deeper, cool waters of the pond.³




Figure 4. 2021 Dissolved Oxygen and Temperature Profiles, Station 1

³ Maine Department of Inland Fisheries & Wildlife. Lake Survey Maps – David Pond. Accessed online: https://www.maine.gov/ifw/docs/lake-survey-maps/kennebec/david_pond.pdf.

Water surface temperatures through the 2021 monitoring season ranged from 20.0 C (68 F) to 26.8 C (80 F) with an average surface water temperature of 23.4 C (74 F) between June and September. Continued collection of bi-weekly DO and temperature profiles will identify trends and changes occurring in David Pond in order to better understand variations in thermal stratification and the extent and severity of the low DO and anoxic zones throughout the monitoring season.

Total Phosphorus (TP)

Phosphorus is the nutrient that most influences the growth of algae in lakes. Because its natural occurrence in lakes is very small, phosphorus “limits” the growth of algae in lake ecosystems. Small increases in phosphorus in lake water can cause substantial

increases in algal growth, hindering lake health as well as the economic, recreational, and aesthetic value of the lake. Tracking in-lake phosphorus levels over time is another way of monitoring change in lake water quality trends.

Total Phosphorus (ppb)	
2021 TP Average	7
Historical TP Average	7
Maine Lakes TP Average	12

Nine (9) samples were collected by 30 Mile staff this year and analyzed for Total Phosphorus (TP). Samples were collected monthly between June and September. Five (5) of the phosphorus samples were collected from the top layer of David Pond using an integrated core sampler and are referred to as “epilimnetic core samples”. Laboratory results for epilimnetic core samples collected in 2021 ranged from 5 ppb to 8 ppb with an annual average of 7 ppb.

Generally speaking, in-lake phosphorus concentrations (epilimnetic samples) less than 10-12 ppb are ideal. Lakes with in-lake phosphorus concentrations of 13 ppb or more are able to sustain algal blooms, and blooms become frequent as in-lake average concentrations approach 20 ppb. Historically, the annual average in-lake phosphorus concentration in David Pond has ranged from 6 ppb (1982, 2016, and 2017) to 10 ppb (2020) with a historical average annual concentration of 7 ppb (Figure 5). TP data for David Pond is also quite limited, with data collected only 8 years throughout the ~40-year monitoring record starting in 1982.

In 2021, four (4) TP samples were collected from the bottom of David Pond using a Kemmerer grab sampler; this type of sample is known as a “bottom grab”. Bottom grabs are collected when anoxia is encountered anywhere in the dissolved oxygen profile, and help us determine if there is active phosphorus release from bottom sediments exposed to anoxic conditions. Laboratory results for bottom grab samples collected in 2021 were 15 ppb (June 18th), 20 ppb (July 27th and August 24th), and 12 ppb (September 22nd) with a 2021 annual average of 17 ppb.

Historically, bottom grab samples were collected in 7 years throughout the ~40-year monitoring record. The historical average annual bottom grab TP concentration ranges from 12 ppb (2013) to 30 ppb (2018) with an overall historical average of 21 ppb.




Figure 5. Annual Average Total Phosphorus data (epilimnetic core samples) collected 1982-2021, Station 1.

Chlorophyll (Chl-a)

Chlorophyll is found in plants (including algae), and is used to convert sunlight into energy. Measuring the concentration of Chlorophyll in lake water helps us estimate the algae population in the lake.

Chl-a (ppb)	
2021 Chl-a Average	4.4
2021 Peak Chl-a	5.0
Historical Chl-a Average	3.1
Maine Lakes Chl-a Average	5.4




Figure 6. Annual average and peak chlorophyll-a, station 1, 1982-2021

Chlorophyll was measured five (5) times in 2021. Results ranged from 3 ppb to 5 ppb, with a 2021 annual average of 4.7 ppb. Historical Chl-a data collected during 7 years between 1982-2021 has ranged from 1.0 ppb (2018) to 7 ppb (2013) with a historical average of 3.5 ppb (Figure 6).

Discussion

2021 was 30 Mile's sixth year of monitoring David Pond. Historical data presented in this report includes all monitoring data collected through 2018, submitted by volunteer monitors, 30 Mile staff, and other state agencies, that has undergone a thorough QA/QC process at Maine DEP. 2019, 2020, and 2021 data included in this report is data collected by 30 Mile only. Annual averages will be updated in next year's water quality report if Maine DEP has published the full dataset by the time of the report.

Five years of consecutive data collection for any given parameter will provide the baseline condition for water quality. 10 years of consecutive data collection is needed to meet the minimum data thresholds for determining trends over time. 30 Mile's monitoring program will continue to develop a robust dataset that can help our community identify and address water quality concerns in David Pond.

Near real-time data for David Pond's clarity (Secchi depth), and dissolved oxygen and temperature profiles can be found online at <https://30mileriver.org/david-pond/>, along with a link to the historical dataset and depth map.

Next Steps

1. Continue **bi-weekly baseline monitoring** between May and October each year to monitor seasonal and annual variability across all parameters, and better document changes and trends over time.
2. Continue to **deliver LakeSmart programming** on David Pond, providing education to shorefront property owners about polluted stormwater runoff, phosphorus, and the affects that watershed development can have on lake water quality.
3. Work with 30 Mile to **review the list of priority sites identified during the 2011 watershed survey** and determine next steps to address remaining sites through LakeSmart and 30 Mile's YCC and Technical Assistance programs.